arXiv:math/9307206v1 [math.CA] 9 Jul 1993

The q-Harmonic Oscillator and an Analog of the Charlier polynomials

R. Askey† and S. K. Suslov‡

Abstract. A model of a q-harmonic oscillator based on q-Charlier polynomials of Al-Salam and Carlitz is discussed. Simple explicit realization of q-creation and q-annihilation operators, q-coherent states and an analog of the Fourier transformation are found. A connection of the kernel of this transform with biorthogonal rational functions is observed.

Models of q-harmonic oscillators are being developed in connection with quantum groups and their various applications (see, for example, Refs. [1–5]). The q-analogs of boson operators were introduced explicitly in Refs. [1,3] and [5], where the corresponding wave functions were found in terms of the continuous q-Hermite polynomials of Rogers [6,7] and in terms of the Stieltjes–Wigert polynomials [8,9], respectively. Here we introduce one more explicit realization of q-creation and q-annihilation operators with the aid of q-Charlier polynomials of Al-Salam and Carlitz [10].

The q-orthogonal polynomials V_n^a studied by Al-Salam and Carlitz may be considered as a *q-version of the Charlier polynomials* $c_n^\mu(s)$ (see, for example, [11,12]). To emphasize this analogy we use the notation $c_n^\mu(x \mid q)$ for the Al-Salam and Carlitz polynomials. In our notation they can be defined by the three-term recurrence relation

$$\mu q^{-n-1} c_{n+1}^\mu(x \mid q) + (1-q^n) q^{-n} c_{n-1}^\mu(x \mid q) = \left((\mu+q)q^{-n-1} - x\right) c_n^\mu(x \mid q)\,, \tag{1}$$

$c_0^\mu(x \mid q) = 1\,, c_1^\mu(x \mid q) = \mu^{-1}(\mu + q - qx)\,.$ These polynomials are orthogonal

$$\sum_{s=0}^{\infty} c_m^\mu(q^{-s} \mid q)\, c_n^\mu(q^{-s} \mid q)\, \rho(s) q^{-s} = \frac{(q;q)_n}{\mu^n}\delta_{mn} \tag{2}$$

with respect to a positive measure

$$\rho(s) = (\mu;q)_\infty \frac{\mu^s q^{s^2}}{(q,\mu;q)_s};\ 0 < \mu, q < 1\,, \tag{3}$$

where the usual notations (see [13]) are

$$\begin{aligned} (a;q)_n &= \prod_{k=0}^{n-1}(1-aq^k)\,, \\ (a,b;q)_n &= (a;q)_n (b;q)_n\,, \\ (a;q)_\infty &= \lim_{n\to\infty}(a;q)_n\,. \end{aligned} \tag{4}$$

---

† Department of Mathematics, University of Wisconsin, Madison, WI 53706, USA
‡ Russian Scientific Center “Kurchatov Institute”, Moscow 123182, Russia

The weight function (3) is a solution of the Pearson equation $\Delta(\sigma\rho) = \rho\tau\nabla x_1$ (for details, see [12,14]) with $x(s) = q^{-s}$, $\sigma(s) = (1-q^{-s})(\mu - q^{1-s})$, $\sigma(s) + \tau(s)\nabla x_1(s) = \mu$. The explicit form of the polynomials $c_n^\mu(x \mid q)$ is

$$c_n^\mu(x \mid q) = {}_2\varphi_0(q^{-n}, x; q, q^{1+n}/\mu),\ x = q^{-s}\,. \tag{5}$$

For the definition of the basic hypergeometric function ${}_2\varphi_0$, see [13]. In the limit $q \to 1$ it easy to see from (1) or (5) that

$$\lim_{q\to 1} c_n^{(1-q)\mu}(q^{-s} \mid q) = {}_2F_0(-n, -s; -1/\mu) = c_n^\mu(s)\,. \tag{6}$$

This justifies our notation for the Al-Salam and Carlitz polynomials.

The polynomials $c_n^\mu(x \mid q)$ give us the possibility to introduce a new model of a q-oscillator. We can define a *q-version of the wave functions* of harmonic oscillator as

$$\psi_n(s) = d_n^{-1} q^{-\frac{s}{2}} \rho^{\frac{1}{2}}(s)\, c_n^\mu(q^{-s} \mid q)\,, \tag{7}$$

where $d_n^2 = (q;q)_n/\mu^n$. These q-wave functions satisfy the orthogonality relation

$$\sum_{s=0}^{\infty} \psi_n(s)\, \psi_m(s) = \delta_{nm}\,. \tag{8}$$

The *q-annihilation* a and *q-creation* a^+ operators have the following explicit form

$$\begin{aligned} a =&(1-q)^{-\frac{1}{2}} \left[\mu^{\frac{1}{2}} q^s - \sqrt{(1-q^{s+1})(1-\mu q^s)}\, e^{\partial_s}\right], \\ a^+ =&(1-q)^{-\frac{1}{2}} \left[\mu^{\frac{1}{2}} q^s - e^{-\partial_s}\sqrt{(1-q^{s+1})(1-\mu q^s)}\right], \end{aligned} \tag{9}$$

where $\partial_s \equiv \dfrac{d}{ds}$, $e^{\alpha\partial_s} f(s) = f(s+\alpha)$. They satisfy the q-commutational rule

$$aa^+ - qa^+a = 1 \tag{10}$$

and act on the q-wave functions defined in (7) by

$$a\psi_n = e_n^{\frac{1}{2}} \psi_{n-1},\ a^+\psi_n = e_{n+1}^{\frac{1}{2}} \psi_{n+1}\,, \tag{11}$$

where

$$e_n = \frac{1-q^n}{1-q}\,.$$

In this model of the q-oscillator equations (11) are equivalent to difference-differentiation formulas

$$\begin{aligned} \mu q^s \Delta c_n^\mu(x \mid q) &= (q^n - 1)\, c_{n-1}^\mu(x \mid q)\,, \\ q^s \nabla[\rho(s) c_n^\mu(x \mid q)] &= \rho(s)\, c_{n+1}^\mu(x \mid q)\,, \end{aligned}$$

respectively. Here $\Delta f(s) = \nabla f(s+1) = f(s+1) - f(s)$ and $x = q^{-s}$. In view of (6) the functions $\psi_n(s)$ converge in the limit $q \to 1^-$ to the wave functions of the discrete model of the linear harmonic oscillator considered in [15].

The *q-Hamiltonian* $H = a^+a$ acts on the wave functions (7) as

$$H\psi_n = e_n\psi_n \tag{12}$$

and has the following explicit form

$$\begin{aligned} H = (1-q)^{-1}\big[&\mu q^{2s} + (1-q^s)(1-\mu q^{s-1}) - \\ &\mu^{\frac{1}{2}} q^s \sqrt{(1-q^{s+1})(1-\mu q^s)}e^{\partial_s} - \mu^{\frac{1}{2}} q^{s-1}\sqrt{(1-q^s)(1-\mu q^{s-1}}e^{-\partial_s}\big]\,. \end{aligned} \tag{13}$$

By factorizing the Hamiltonian (or the difference equation for the Al-Salam and Carlitz polynomials) we arrive at the explicit form (9) for the q-boson operators.

Since $a^+a = H$, the relation (10) can be written in the equivalent form

$$[a, a^+] = 1 - (1-q)H \equiv q^N\,. \tag{14}$$

The operator

$$N = \frac{1}{\log q}\log[1 - (1-q)H] \tag{15}$$

can be considered as the *number operator*, since

$$[a, N] = a, \qquad [N, a^+] = a^+\,. \tag{16}$$

From these relations one can obtain the equations (11) and the spectrum (12) of the q-Hamiltonian in abstract form. The q-wave functions are

$$\psi_n(s) = c_n\left(a^+\right)^n\psi_0(s)\,, \qquad a\,\psi_0(s) = 0\,,$$

where $c_n = (e_n!)^{-1/2}$ and $e_n! = e_1e_2\ldots e_n$.

For the model of the q-oscillator under discussion we can construct explicitly *q-coherent states* and an analog of the Fourier transformation. For the coherent states $\mid \alpha\rangle$ defined by

$$\begin{aligned} a \mid \alpha\rangle &= \alpha \mid \alpha\rangle\,, \qquad \langle\alpha|\alpha\rangle = 1\,, \\ \mid \alpha\rangle &= f_\alpha \sum_{n=0}^{\infty} \frac{\alpha^n\psi_n(s)}{(e_n!)^{1/2}}\,, \\ f_\alpha &= \left((1-q) \mid \alpha|^2; q\right)_\infty^{\frac{1}{2}}\,, \quad (1-q) \mid \alpha|^2 < 1 \end{aligned} \tag{17}$$

we can write

$$\mid \alpha\rangle = f_\alpha\left(\rho q^{-s}\right)^{\frac{1}{2}} \sum_{n=0}^{\infty} \frac{t^n}{(q;q)_n}\, c_n^\mu(x \mid q)\,, t = \alpha\mu^{\frac{1}{2}}(1-q)^{\frac{1}{2}}\,. \tag{18}$$

With the aid of the generating function [10]

$$\sum_{n=0}^{\infty} \frac{t^n}{(q;q)_n}\, c_n^{\mu}(x \mid q) = \frac{(qtx/\mu;q)_\infty}{(t\,,qt/\mu;q)_\infty}\,,\ |t|<1,\ |qt/\mu|<1 \tag{19}$$

we arrive at the following *explicit form* for the q-coherent states

$$\mid \alpha\rangle = f_\alpha\left(\rho q^{-s}\right)^{\frac{1}{2}} \frac{\left(\alpha(1-q)^{1/2}\mu^{-1/2}q^{1-s};q\right)_\infty}{\left(\alpha(1-q)^{1/2}\mu^{1/2}, \alpha q(1-q)^{1/2}\mu^{-1/2};q\right)_\infty}\,, \tag{20}$$

where $\rho = (q;q)_\infty^{-1}(q^{s+1}\,,\mu q^s;q)_\infty \mu^s q^{s^2}$. These coherent states are not orthogonal

$$\langle\alpha \mid \beta\rangle = \frac{\left((1-q)\mid\alpha\mid^2, (1-q)\mid\beta\mid^2;q\right)_\infty^{1/2}}{\left((1-q)\alpha^*\beta\,;q\right)_\infty}\,,$$

where $*$ denotes the complex conjugate.

To define an analog of the *Fourier transform* we can consider, following Wiener's approach to the classical Fourier transform [16] (see also [17,18]), the kernel of the form

$$\begin{aligned} K_t(s,p) &= \sum_{n=0}^{\infty} t^n \psi_n(s)\,\psi_n(p) \\ &= \left(\rho(s)\rho(p)q^{-s-p}\right)^{\frac{1}{2}} \sum_{n=0}^{\infty} \frac{(\mu t)^n}{(q;q)_n}\, c_n^{\mu}(q^{-s}\mid q)\, c_n^{\mu}(q^{-p}\mid q)\,. \end{aligned} \tag{21}$$

The series can be summed with the aid of the bilinear generating function by Al-Salam and Carlitz [10]

$$\begin{aligned} \sum_{n=0}^{\infty} c_n^{\mu_1}(x\mid q)\, c_n^{\mu_2}(y\mid q)\, \frac{t^n}{(q;q)_n} &= \frac{(qtx/\mu_1\,, qty/\mu_2;q)_\infty}{(t\,,qt/\mu_1\,,qt/\mu_2;q)_\infty} \\ &\cdot {}_3\varphi_2\left(\begin{matrix} x\,,y\,,t \\ qtx/\mu_1, qty/\mu_2 \end{matrix}\,;q,\frac{q^2t}{\mu_1\mu_2}\right)\,. \end{aligned} \tag{22}$$

The answer is

$$\begin{aligned} K_t(s,p) &= \left(\rho(s)\rho(p)q^{-s-p}\right)^{\frac{1}{2}} \frac{(tq^{1-s}\,,tq^{1-p}\,;q)_\infty}{(qt\,,qt\,,\mu t;q)_\infty} \\ &\cdot {}_3\varphi_2\left(\begin{matrix} q^{-s}\,,q^{-p}\,,\mu t \\ tq^{1-s}\,,tq^{1-p} \end{matrix}\,;q,\frac{q^2t}{\mu}\right)\,. \end{aligned} \tag{23}$$

The q-wave functions (7) are eigenfunctions of the "discrete q-Fourier transform",

$$i^m\psi_m(s) = \sum_{p=0}^{\infty} K_i(s,p)\,\psi_m(p)\,. \tag{24}$$

The orthogonality relation of the kernel,

$$\sum_{p=0}^{\infty} K_i(s,p)\, K_i^*(s',p) = \delta_{ss'}\,, \tag{25}$$

implies the orthogonality of the rational functions (23). In view of (6) in the limit $q \to 1^-$ we get the "discrete Fourier transform" considered in [17].

Similarly, with the aid of the bilinear generating function (22) and the orthogonality property of the Wall polynomials, which are dual to the polynomials (5), one can obtain the *biorthogonality relation*

$$\sum_{s=0}^{\infty} u_m(s)\, v_n(s)\, \rho(s) q^{-s} = d_n^2 \delta_{mn} \tag{26}$$

with

$$\rho(s) = \frac{\left(\frac{\mu_2}{t_1}, \frac{\mu_2}{t_2}\,; q\right)_s}{(q, \mu_2\,; q)_s}\, \mu_1^s q^s$$

and

$$d_n^2 = \frac{(t_1, t_2\,; q)_\infty}{(\mu_1, \mu_2\,; q)_\infty} \cdot \frac{(q, \mu_1\,; q)_n}{\left(\frac{\mu_1}{t_1}, \frac{\mu_1}{t_2}\,; q\right)_n}\, \mu_2^{-n}$$

for the ${}_3\varphi_2$-*rational functions* of the form

$$u_m(s) = {}_3\varphi_2 \left(\begin{matrix} q^{-m}\,,\ q^{-s}\,,\ t_1 \\ \dfrac{t_1}{\mu_1} q^{1-m}\,,\ \dfrac{t_1}{\mu_2} q^{1-s}\,; q,\ \dfrac{q^2}{t_2} \end{matrix} \right)\,; \tag{27}$$

$$v_n(s) = u_n(s)|_{t_1 \leftrightarrow t_2}\,; \qquad t_1 t_2 = \mu_1 \mu_2\,.$$

These functions are self-dual. They belong to classical biorthogonal rational functions [19,20].

We have considered here the explicit form of q-boson operators which satisfy the commutational rule (10) when $0 < q < 1$. The case $q > 1$ is also interesting. It leads to another family of Al-Salam and Carlitz polynomials.

It is evident that for the models of the q-oscillator under discussion one can readily construct dynamical symmetry group $SU_q(1,1)$ [4] and write an explicit realization for irreducible representations $|j,m\rangle_q = \psi_{j+m}(s)\psi_{j-m}(s')$ of the group$SU_q(2)$ [1,2].

REFERENCES

1. Macfarlane, A.J. (1989) J. Phys. A: Math. Gen., Vol. 22, p. 4581–4588.
2. Biedenharn, L.C. (1989) J. Phys. A: Math. Gen., Vol. 22, p. L873–L878.
3. Atakishiyev, N.M. and Suslov, S.K. (1990) Teor. i Matem. Fiz., Vol. 85, No. 1, p. 64–73.
4. Kulish, P.P. and Damaskinsky, E.V. (1990) J. Phys. A: Math. Gen., Vol. 23, p. L415–L419.
5. Atakishiyev, N.M. and Suslov, S.K. (1991) Teor. i Matem. Fiz., Vol. 87, No. 1, p. 154–156.
6. Rogers, L.J. (1894) Proc. London Math. Soc., Vol. 25, p. 318–343.
7. Askey, R. and Ismail, M.E.H. (1983) In:*Studies in Pure Mathematics* (P. Erdös, ed.), Birkhäuser, Boston, Massachusetts, p. 55–78.
8. Stieltjes, T.J. *Recherches sur les Fractions Continues*, Annales de la Faculté des Sciences de Toulouse, 8 (1894) 122 pp., 9 (1895), 47 pp. Reprinted in *Oeuvres Complétes*, vol. 2.
9. Wigert S. (1923) Arkiv för Matematik, Astronomi och Fysik, Bd. 17, No. 18, p. 1–15.
10. Al-Salam, W.A. and Carlitz, L. (1965) Math. Nachr., Bd. 30, S. 47–61.
11. Chihara, T.S. (1978) *An Introduction to Orthogonal Polynomials.* Gordon and Breach, New York.
12. Nikiforov, A.F., Suslov, S.K., and Uvarov, V.B. (1991) *Classical Orthogonal Polynomials of a Discrete Variable.* Springer-Verlag, Berlin, Heidelberg.
13. Gasper, G. and Rahman, M. (1990) *Basic Hypergeometric Series,* Cambridge Univ. Press, Cambridge.
14. Suslov, S.K. (1989) Russian Math. Surveys, London Math. Soc., Vol. 44, No. 2, p. 227–278.
15. Atakishiyev, N.M. and Suslov, S.K. (1989) A Model of the Harmonic Oscillator on the Lattice. In: *Contemporary Group Analysis: Methods and Applications,* Baku, p. 17–21 [in Russian].
16. Wiener, N. (1933) *The Fourier Integral and Certain of Its Applications.* Cambridge University Press, Cambridge.
17. Askey, R., Atakishiyev, N.M., and Suslov, S.K. Fourier Transformations for Difference Analogs of the Harmonic Oscillator, to appear
18. Askey, R., Atakishiyev, N.M., and Suslov, S.K. An Analog of the Fourier Transformations for a q-Harmonic Oscillator. Preprint No.5611/1, Kurchatov Institute, Moscow, 1993.
19. Wilson, J.A. (1991) SIAM J. Math. Anal., Vol. 22(4), p. 1147–1155.
20. Rahman, M. and Suslov, S.K. Classical Biorthogonal Rational Functions, submitted.